

# HIV CASES AMONG PERSONS RESIDING IN ALABAMA AT DIAGNOSIS

PRELIMINARY 2020 - 4th Quarter (January 1 • December 31)

Race/Ethnicity	% of Newly Diagnosed	% of Prevalent Cases
Black	70.1	63.9
White	22.2	27.1
Hispanic	4.3	3.3
Multi-race	1.5	5.0
Other/Unknown	1.9	0.7
<b>Total</b>	<b>100.0</b>	<b>100.0</b>

Age (Years)	% of Newly Diagnosed	% of Prevalent Cases
<13	0.9	0.2
13-19	5.1	0.5
20-24	21.6	3.5
25-29	20.3	8.7
30-39	24.6	21.4
40-49	13.3	20.9
≥50	14.1	44.9
<b>Total</b>	<b>100.0</b>	<b>100.0</b>

Adult/Adolescent Exposure (≥13 years)	% of Newly Diagnosed	% of Prevalent Cases
Men who have Sex with Men (MSM)	23.3	45.6
Heterosexuals	23.0	29.6
Injection Drug Users (IDU)	1.1	5.0
MSM/IDU	0.9	3.1
Hemophilia/Coagulation Disorder	0.0	0.1
Mother with HIV Infection	0.0	0.6
Transfusion/Transplant Recipient	0.0	0.0
Risk Not Reported/Unknown	51.6	15.9
<b>Total (add pediatric cases to total)</b>	<b>100.0</b>	<b>100.0</b>

Gender	% of Newly Diagnosed	% of Prevalent Cases
Male	76.7	73.2
Female	23.3	26.8
<b>Total (unknowns excluded)</b>	<b>100.0</b>	<b>100.0</b>

Pediatric Exposure (<13 years)	Newly Diagnosed		Prevalent Cases	
	Cases	% of Total	Cases	% of Total
Mother with HIV Infection	4	0	21	77.8
Hemophilia/Coagulation Disorder	0	0	0	0.0
Transfusion/Transplant Recipient	0	0	0	0.0
Risk Not Reported/Unknown	1	0	6	22.2
<b>Total</b>	<b>5</b>	<b>0</b>	<b>27</b>	<b>100.0</b>

Public Health District	Newly Diagnosed		Prevalent Cases	
	Cases	% of Total	Cases	% of Total
Northern	59	12.1	1676	11.3
East Central	107	22.0	2934	19.8
West Central	52	10.7	1050	7.1
Jefferson	100	20.6	3920	26.5
Northeastern	36	7.4	1336	9.0
Southeastern	42	8.6	1084	7.3
Southwestern	23	4.7	781	5.3
Mobile	67	13.8	2010	13.6
<b>Total (does not include unknown)</b>	<b>486</b>	<b>100.0</b>	<b>14791</b>	<b>100.0</b>

**\*Note: Statistics should be interpreted WITH CAUTION as not all reported cases have been entered into the HIV Surveillance database. Effective October 1, 2017, Public Health Areas have been redistributed as eight Public Health Districts. Unknown cases are only accounted for in state total.**  
 Newly diagnosed HIV includes newly diagnosed HIV infections during the year of interest.  
 Prevalent HIV includes all persons living with HIV as of December 28, 2020. Cumulative HIV includes all diagnosed HIV (living and deceased) as of December 28, 2020.  
 Totals include unknown case counts. Females with no risk factor reported are reclassified as heterosexual exposure.  
 Age among newly diagnosed and cumulative cases is age at diagnosis. Prevalent age is current age among cases living as of December 28, 2020.  
 Public Health District represents residence at diagnosis among newly diagnosed and cumulative cases and current residence among prevalent cases.