Alabama Prenatal STD Testing

Test (All tests must be FDA approved)	Initial Prenatal Visit	Third Trimester	Labor and Delivery (L&D)
Syphilis	All pregnant persons.	All pregnant persons at 28-32 weeks gestation, regardless of risk factors.	All pregnant persons.
HIV	All pregnant persons not previously confirmed as HIV infected.	All pregnant persons at 28-32 weeks gestation, regardless of risk factors unless previously confirmed as HIV infected.	All pregnant persons unless already confirmed as HIV infected.
Chlamydia	All pregnant persons.	All pregnant persons at 36 weeks gestation if the initial test was positive, have signs and symptoms, or at high risk of infections.	
Gonorrhea	All pregnant persons.	All pregnant persons at 36 weeks gestation if the initial test was positive, have signs and symptoms, or at high risk of infections.	
HBV	All pregnant persons.		All pregnant persons if no prior testing or have signs and symptoms of hepatitis.
НСУ	All pregnant persons.		

No Prenatal Care – Patient Presents at Delivery

Test	Labor and Delivery		
Syphilis	All pregnant persons.		
HIV	All pregnant persons unless already confirmed to be infected with HIV infection.		
Chlamydia	All pregnant persons.		
Gonorrhea	All pregnant persons.		
HBV	All pregnant persons.		
HCV	All pregnant persons.		