

Promising actions for safer opioid prescribing



Problem: High prescribing
Solution: Safer prescribing practices



Problem:
Too many prescriptions



In 2015, the amount of opioids prescribed was enough for every American to be medicated **around the clock for 3 weeks.**

(640 MME per person, which equals 5 mg of hydrocodone every 4 hours)



Solution:
Fewer prescriptions


Use opioids **only** when benefits are likely to outweigh risks. Options other than opioids include:

- Pain medicines like acetaminophen, ibuprofen, and naproxen
- Physical therapy and exercise
- Cognitive behavioral therapy

Therapies that don't involve opioids may work better and have fewer risks and side effects.



Problem:
Too many days



Average days supply per prescription increased from 2006 to 2015.

Even at low doses, taking an opioid for more than 3 months increases the risk of addiction by **15 times.**




Solution:
Fewer days

For acute pain, prescriptions should only be for the expected duration of pain severe enough to need opioids. **Three days or less** is often enough; more than seven days is rarely needed.

If continuing opioids, ask whether benefits continue to outweigh risks. If not, use other treatments and taper opioids gradually.



Problem:
Too high a dose



Average daily MME per prescription declined both nationwide and in most counties, but it is still too high.

A dose of 50 MME or more per day **doubles** the risk of opioid overdose death, compared to 20 MME or less per day. At 90 MME or more, the risk increases **10 times.**



Solution:
Lower doses

Use the lowest effective dose of immediate-release opioids when starting, and reassess benefits and risks when considering dose increases.

Avoid a daily dose of 90 MME or more. If already taking high doses, offer the opportunity to gradually taper to safer doses.

For more recommendations when considering opioids for chronic pain outside of end-of-life care, see **The CDC Guideline for Prescribing Opioids for Chronic Pain.** The Guideline can also be used to inform health systems, states, and insurers to ensure appropriate prescribing and improve care for all people.

www.cdc.gov/drugoverdose/prescribing/guideline.html

What Can Be Done?



The Federal Government is

- Educating healthcare providers and the public about pain management, addiction, and opioid overdose and providing guidance on safe and effective pain management.
- Equipping states with resources to implement and evaluate safe prescribing practices.
- Improving access to addiction treatment and recovery services.
- Increasing access to overdose-reversing drugs, such as naloxone.
- Tracking opioid-related trends to better understand and respond to the epidemic.
- Supporting cutting-edge research about pain management and addiction.

States can

- Maximize prescription drug monitoring programs (PDMPs) by using near real-time data reporting, integrating with electronic health records, and promoting routine provider use.
- Implement and evaluate programs to improve prescribing practices.
- Use data to identify and address high-risk prescribing.
- Enhance the use of prescribing guidelines based on the best available science.
- Increase access to medication-assisted treatment for addiction and naloxone for opioid overdose.

Health insurers can

- Refer to the Guideline in setting up prescription claims review programs to identify and address improper prescribing and use of opioids.
- Increase coverage for other proven treatments to reduce pain, such as physical therapy and non-opioid pain medicines.

- Cover clinicians' time when they are conducting activities that improve quality and safety of pain management and addressing addiction. These can include patient counseling, coordination of care, and checking the patient's prescription history in the PDMP.
- Reduce barriers (such as prior authorization) to use of nonopioid pain medications and medication-assisted treatment for addiction.

Healthcare providers can:

- Follow the *CDC Guideline for Prescribing Opioids for Chronic Pain*, which includes recommendations such as:
 - ▶ Use opioids only when benefits are likely to outweigh risks.
 - ▶ Start with the lowest effective dose of immediate-release opioids. For acute pain, prescribe only the number of days that the pain is expected to be severe enough to require opioids.
 - ▶ Reassess benefits and risks if considering dose increases.
- Use state-based PDMPs which help identify patients at risk of addiction or overdose.

Everyone can

- Consider non-opioid options for pain management.
- Store prescription opioids in a secure place, out of reach of others (including children, family, friends, and visitors).
- Dispose of medications properly as soon as the course of treatment is done. For more information, visit FDA at: www.fda.gov/Drugs/ResourcesForYou.
- Get help if you're having trouble controlling your opioid use, SAMHSA's National Helpline: [1-800-662-HELP](tel:1-800-662-HELP).

1-800-CDC-INFO (232-4636)

TTY: 1-888-232-6348

www.cdc.gov

Centers for Disease Control and Prevention

1600 Clifton Road NE, Atlanta, GA 30329

Publication date: 7/6/2017