

**Pharmacology Update: Interaction of Oral Contraceptives and  
Seizure Medications to Drugs and Herbal Remedies**

Satellite Webcast

Social Work CEU Quiz

March 4, 2015

1. A patient taking an OC and a protease inhibitor should always use a barrier method of contraception, regardless of the interaction level:  
A) True  
B) False
2. A medication listed as a level 2 interaction should never be given to a patient receiving oral contraceptives  
A) True  
B) False
3. The breast cancer prophylaxis medication, Tamoxifen (Nolvadex) is:  
A) Contraindicate in most, but not all patients  
B) Oral contraceptives may aggravate conditions being treated by Tamoxifen  
C) All of the above
4. Topiramate (Topamax) level 2 interactions include:  
A) An increase clearance of Ethinyl Estradiol in oral and non-oral contraceptives  
B) Ethanol alters pattern of Topiramate release from extended-release capsules  
C) Decrease of Dolutegravir levels  
D) All of the above
5. Phenytoin (Dilantin) has over \_\_\_\_\_ drugs listed as having the potential for a level 4 interaction:  
A) 10  
B) 15  
C) 20  
D) 30

Name: \_\_\_\_\_

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Phone #: \_\_\_\_\_ Email: \_\_\_\_\_