## Early Hearing Detection and Intervention (EHDI) and the Newborn Screening Program Social Work CEU Quiz

July 12, 2016

(Select only the one best answer)

1. \_\_\_\_\_ is more prevalent than all other newborn screening disorders

|       | A) Endocrine disorders  |
|-------|---|
|       | B) Hearing loss   |
|       | C) Metabolic disorders  |
|       | D) Cystic Fibrosis  |
| 2.    | A hearing screen is a noninvasive, automated test to detect potential hearing loss:         |
|       | A) True   |
|       | B) False  |
| 3.    | Types of hearing loss include:  |
|       | A) Sensorineural Hearing Loss   |
|       | B) Conductive Hearing Loss  |
|       | C) Auditory Neuropathy Spectrum Disorder  |
|       | D) All of the above   |
| 4.    | Neonatal Intensive Care Unit (NICU) stays of less than 5 days are a risk factor for hearing |
|       | loss.   |
|       | A) True   |
|       | B) False  |
| 5.    | The most common type of permanent hearing loss is:  |
|       | A) Sensorineural Hearing Loss   |
|       | B) Conductive Hearing Loss  |
|       | C) Auditory Neuropathy Spectrum Disorder  |
|       | D) All of the above   |
| Name  |   |
| Addre | ss: State: Zip:   |
| Phone | #: Email:   |
|       |   |