OASIS ITEM

(M1308) Current Number of Unhealed (non epithelialized) Pressure Ulcers at Each Stage: (Enter "0" if none; excludes Stage I pressure ulcers)

		COLUMN 1	COLUMN 2
		Complete at	Complete at
		SOC/ROC/FU & D/C	FU & D/C
Stage description – unhealed pressure ulcers		Number Currently Present	Number of those listed in Column 1 that were present on admission (most recent SOC / ROC)
a.	Stage II: Partial thickness loss of dermis presenting as a shallow open ulcer with red pink wound bed, without slough. May also present as an intact or open/ruptured serum-filled blister.		
b.	Stage III: Full thickness tissue loss. Subcutaneous fat may be visible but bone, tendon, or muscles are not exposed. Slough may be present but does not obscure the depth of tissue loss. May include undermining and tunneling.		
C.	Stage IV: Full thickness tissue loss with visible bone, tendon, or muscle. Slough or eschar may be present on some parts of the wound bed. Often includes undermining and tunneling.		
d.1	Unstageable: Known or likely but not stageable due to non-removable dressing or device		
d.2	Unstageable: Known or likely but not stageable due to coverage of wound bed by slough and/or eschar.		
d.3	Unstageable: Suspected deep tissue injury in evolution.		

ITEM INTENT

Identifies the number of Stage II or higher pressure ulcers at each stage present at the time of assessment. Stage I pressure ulcers are <u>not</u> reported in this item.

TIME POINTS ITEM(S) COMPLETED

Start of care - Column 1

Resumption of care - Column 1

Follow-up - Columns 1 and 2

Discharge from agency - not to inpatient facility - Columns 1 and 2