



INFANT MORTALITY ALABAMA 2017


**ALABAMA DEPARTMENT OF PUBLIC HEALTH
CENTER FOR HEALTH STATISTICS**




INFANT MORTALITY RATES ALABAMA, 2008-2017



NUMBER OF INFANT DEATHS ALABAMA, 2008-2017




NUMBER OF BIRTHS ALABAMA, 2008-2017




INFANT MORTALITY BY COUNTY OF RESIDENCE

ALABAMA, 2017


Number




Rate (Per 1,000 Live Births)




INFANT MORTALITY RATES ALABAMA, 2008-2017



THREE-YEAR INFANT MORTALITY RATES ALABAMA, 1995-1997 to 2015-2017




INFANT MORTALITY RATES ALABAMA AND UNITED STATES¹, 1970-2017




¹ Most current national data is for year 2016.


NUMBER OF BIRTHS TO TEENAGERS (AGES 10-19) ALABAMA, 2008-2017



PERCENT OF BIRTHS TO TEENAGERS (AGES 10-19) ALABAMA, 2008-2017




PERCENT OF BIRTHS TO TEENAGERS BY RACE (AGES 10-19) ALABAMA, 2016 vs. 2017



PERCENT OF BIRTHS LESS THAN 37 WEEKS GESTATION ALABAMA, 2008-2017




PERCENT OF BIRTHS AT 37- 40 WEEKS GESTATION ALABAMA, 2011-2017




PERCENT OF LOW WEIGHT BIRTHS ALABAMA, 2008-2017



NUMBER OF PRETERM BIRTHS BY BIRTH WEIGHT ALABAMA, 2016 vs. 2017



NUMBER OF INFANT DEATHS WITH PRETERM BIRTHS BY BIRTH WEIGHT ALABAMA, 2016 vs. 2017



PERCENT OF BIRTHS WITH A BIRTH INTERVAL LESS THAN 2 YEARS, ALABAMA, 2008-2017




PERCENT OF BIRTHS WITH ADEQUATE PRENATAL CARE¹ ALABAMA, 2008-2017




¹ Adequacy of care included both adequate plus and adequate as measured by the Kotelchuck index. Percentage calculation excludes unknown status.

PERCENT OF BIRTHS WITH NO PRENATAL CARE ALABAMA, 2008-2017




PERCENT OF BIRTHS WITH MATERNAL SMOKING ALABAMA, 2008-2017



* Smoking question changed from “yes” or “no” during pregnancy to an indication for each trimester in 2014. For comparison, smoking in any trimester equaled smoking during pregnancy.

PERCENT OF BIRTHS WITH MATERNAL SMOKING BY AGE OF MOTHER ALABAMA, 2008-2017



* Smoking question changed from “yes” or “no” during pregnancy to an indication for each trimester in 2014. For comparison, smoking in any trimester equaled smoking during pregnancy.

LEADING CAUSES OF INFANT DEATHS ALABAMA, 2017

- Top three leading causes of infant deaths
 - Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)
 - Disorders related to short gestation and low birth weight, not elsewhere classified (P07)
 - Sudden infant death syndrome (R95)
- Three leading causes accounted for 43.4% of infant deaths.
- Top three leading causes of infant deaths were same for US in 2016 and accounted for 44.2% of infant deaths.

LEADING CAUSES OF INFANT DEATHS ALABAMA, 2015-2017

2015	White	Black	Other	Total
Q00-Q99	43	42	1	86
P07	27	42	0	69
R95	20	41	0	61
Subtotal	90	125	1	216
Total Deaths	206	280	8	494
Top 3				43.7%

2016	White	Black	Other	Total
Q00-Q99	55	35	2	92
P07	28	60	1	89
R95	27	22	2	51
Subtotal	110	117	5	232
Total Deaths	250	275	12	537
Top 3				47.0%

2017	White	Black	Other	Total
Q00-Q99	53	33	5	91
P07	19	39	1	59
R95	23	16	0	39
Subtotal	95	88	6	189
Total Deaths	213	207	15	435
Top 3				43.4%