

Tdap	HPV	MenACWY	Flu
<p><b>Tetanus &amp; diphtheria toxoids &amp; acellular pertussis vaccine (Tdap)</b>  <a href="#">Summary of ACIP</a>  <a href="#">Recommendations for Tdap</a></p> <ul style="list-style-type: none"> <li>• Administer Tdap at age 11-12, as well as at age 13-18 if they have not yet received Tdap, followed by Td booster dose every 10 years.</li> <li>• Those age 7-10 not fully immunized, never vaccinated, or have unknown status against pertussis should receive single dose of Tdap. Refer to the catch-up schedule if additional doses of Td-containing vaccine are needed.</li> <li>• Tdap can be administered regardless of interval since the last Td-containing vaccine.</li> <li>• Administer 1 dose of Tdap vaccine to pregnant adolescents during each pregnancy (preferred during 27 through 36 weeks gestation) regardless of time since prior Td or Tdap vaccination.</li> </ul>	<p><b>Human papillomavirus vaccine (HPV)</b>  <a href="#">Summary of ACIP</a>  <a href="#">Recommendations for HPV</a></p> <ul style="list-style-type: none"> <li>• HPV4 or HPV2 is recommended for the prevention of cervical precancers and cancers in females.</li> <li>• HPV4 is recommended for prevention of cervical, vaginal, vulvar, anal precancers and cancers as well as genital warts in females.</li> <li>• HPV4 is recommended for prevention of anal precancers and cancers, as well as genital warts in males.</li> <li>• HPV4 or HPV2 is recommended at age 11-12 years but can be given as early as age 9.</li> <li>• HPV vaccine is a 3-dose series. Administer the second dose 1 to 2 months after the first dose and the third dose 6 months after the first dose (at least 24 weeks after the first dose).</li> </ul>	<p><b>Meningococcal conjugate vaccine, quadrivalent (MenACWY)</b>  <a href="#">Summary of ACIP</a>  <a href="#">Recommendations for MenACWY</a></p> <ul style="list-style-type: none"> <li>• Administer MCV4 at age 11-12 with a booster dose at age 16 years.</li> <li>• If the first dose of MCV4 is given after the 16th birthday, no booster is recommended.</li> <li>• Administer 1 dose at age 13-18 if not previously vaccinated.</li> <li>• Persons who received their first dose at age 13-15 should receive a booster dose at age 16-18 years.</li> <li>• Administer 1 dose to previously unvaccinated college freshmen living in a dormitory or military recruits living in barracks.</li> </ul>	<p><b>Influenza vaccine (seasonal)</b>  <a href="#">Summary of ACIP</a>  <a href="#">Recommendations for Flu</a></p> <ul style="list-style-type: none"> <li>• Administer influenza vaccine annually to all children beginning at age 6 months — including preteens and teens.</li> <li>• For healthy non-pregnant persons age 7-18 (i.e., those who do not have underlying medical conditions that predispose them to influenza complications), either LAIV or IIV may be used.</li> </ul>