## ALABAMA PRENATAL STD TESTING



<b>Test</b> (All test must be FDA approved)	Initial Prenatal Visit	Third Trimester	Labor and Delivery (L&D)
Syphilis	All pregnant persons.	All pregnant persons at 28-32 weeks gestation.	All pregnant persons.
HIV	All pregnant persons not previously confirmed as HIV infected.	All pregnant persons at 28-32 weeks gestation, regardless of risk factors unless previously confirmed as HIV infected.	All pregnant persons unless already confirmed as HIV infected.
Chlamydia	All pregnant persons.	All pregnant persons at 36 weeks gestation if: • The initial test was positive, • Signs or symptoms present, or • At high risk of infections	Only if no prenatal care.
Gonorrhea	All pregnant persons.	All pregnant persons at 36 weeks gestation if: • The initial test was positive, • Signs or symptoms present, or • At high risk of infections	Only if no prenatal care.
HBV	All pregnant persons.		All pregnant persons.
HCV	All pregnant persons.		Only if no prenatal care.

## No Prenatal Care -Patient Presents at Delivery

Test	Labor and Delivery		
Syphilis	All pregnant persons.		
HIV	All pregnant persons unless already confirmed to be infected with HIV infection.		
Chlamydia	All pregnant persons.		
Gonorrhea	All pregnant persons.		
HBV	All pregnant persons.		
HCV	All pregnant persons.		